

EXTREMES OF INDEPENDENT GAUSSIAN PROCESSES

ZAKHAR KABLUCHKO

ABSTRACT. For every $n \in \mathbb{N}$, let $X_{1n}, \dots, X_{nn}$ be independent copies of a zero-mean Gaussian process $X_n = \{X_n(t), t \in T\}$. We describe all processes which can be obtained as limits, as $n \rightarrow \infty$, of the process $a_n(M_n - b_n)$, where $M_n(t) = \max_{i=1, \dots, n} X_{in}(t)$ and a_n, b_n are normalizing constants. We also provide an analogous characterization for the limits of the process $a_n L_n$, where $L_n(t) = \min_{i=1, \dots, n} |X_{in}(t)|$.

1. INTRODUCTION

Suppose that we are given a large number n of independent copies of some Gaussian process defined on an arbitrary set T . Let M_n be the maximum of these processes, taken pointwise. The aim of this paper is to describe the class of processes which can be obtained as limits, as $n \rightarrow \infty$ and after suitable normalization, of the process M_n . To state our problem more precise, let $X_{1n}, \dots, X_{nn}$ be n independent copies of a zero-mean Gaussian process $\{X_n(t), t \in T\}$ with

$$(1) \quad r_n(t_1, t_2) := \mathbb{E}[X_n(t_1)X_n(t_2)], \quad \sigma_n^2(t) := r_n(t, t) > 0.$$

We define a process $\{M_n(t), t \in T\}$ by

$$(2) \quad M_n(t) = \max_{i=1, \dots, n} X_{in}(t).$$

Problem 1. Describe all sequences X_n and all sequences of constants a_n, b_n for which the process $a_n(M_n - b_n)$ converges as $n \rightarrow \infty$ to some nontrivial limit.

Throughout, convergence of stochastic processes is understood as the weak convergence of their finite-dimensional distributions.

Hüsler and Reiss [5] proved that M_n converges as $n \rightarrow \infty$ to a nontrivial limiting process if the following two conditions are satisfied:

- (1) $\sigma_n(t) = 1$ for all $t \in T$.
- (2) $\lim_{n \rightarrow \infty} \log n \cdot (1 - r_n(t_1, t_2))$ exists in $(0, \infty)$ for all $t_1, t_2 \in T$.

We will complement the work of Hüsler and Reiss [5] by considering processes with *non-unit variance*, providing *necessary and sufficient* conditions for convergence, stressing the role of *negative-definite functions*, and giving a different representation for the limiting processes. This representation is related to a *symmetry property* of certain exponential intensity Poisson point processes with Gaussian increments which is of independent interest.

Extremes of independent *stationary* Gaussian processes on $\mathbb{R}^d$ have been studied by [9]. We will extend some of the results of [9] to the non-stationary case.

2000 *Mathematics Subject Classification.* Primary, 60G70; Secondary, 60G15.

Key words and phrases. Extremes, Gaussian processes, Max-stable processes.

We will also consider a similar problem for *minima* (always understood in the sense of absolute value) of independent Gaussian processes. With the above assumptions, we define a process $\{L_n(t), t \in T\}$ by

$$(3) \quad L_n(t) = \min_{i=1, \dots, n} |X_{in}(t)|.$$

Problem 2. Describe all sequences X_n and all sequences of constants a_n, b_n for which the process $a_n(L_n - b_n)$ converges as $n \rightarrow \infty$ to some nontrivial limit.

The paper is organized as follows. In Section 2 we state Theorem 2.1 which is needed to define the class of limiting processes for both maxima and minima of independent Gaussian processes. We solve Problem 1 and Problem 2 in Section 3 and Section 4. In Section 5 we consider maxima and minima of independent fractional Brownian motions, and construct a new class of α -stable processes. The proof of Theorem 2.1 is given in Section 6.

2. A FAMILY OF GAUSS-POISSON POINT PROCESSES

2.1. Negative-definite kernels. We start by recalling some definitions related to negative-definite kernels, see [2, Chapter 3]. A kernel on a set T is a mapping from $T \times T$ to $\mathbb{R}$. A kernel $\Gamma : T \times T \rightarrow [0, \infty)$ is called *negative definite* if for every $n \in \mathbb{N}$, $t_1, \dots, t_n \in T$ and $a_1, \dots, a_n \in \mathbb{R}$ satisfying $\sum_{i=1}^n a_i = 0$, we have

$$(4) \quad \sum_{i=1}^n \sum_{j=1}^n a_i a_j \Gamma(t_i, t_j) \leq 0.$$

It is well known that $d(t_1, t_2) := \sqrt{\Gamma(t_1, t_2)}$ defines a pseudo-metric on T (which means that $d(t_1, t_1) = 0$, $d(t_1, t_2) = d(t_2, t_1)$, and $d(t_1, t_3) \leq d(t_1, t_2) + d(t_2, t_3)$ for every $t_1, t_2, t_3 \in T$, but note that, in general, $d(t_1, t_2) = 0$ does not imply that $t_1 = t_2$). Further, a pseudo-metric d on T arises in this way for some negative-definite kernel Γ if and only if the pseudo-metric space (T, d) is isometrically embeddable into a Hilbert space.

Example 2.1. If $\{W(t), t \in T\}$ is a zero-mean Gaussian process, then its *incremental variance*

$$\Gamma(t_1, t_2) := \text{Var}(W(t_1) - W(t_2))$$

is a negative-definite kernel on T .

It will be convenient to use the following definition.

Definition 2.1. A kernel $\Gamma : T \times T \rightarrow [0, \infty]$ is called *negative definite in the extended sense* if there is a disjoint decomposition $T = \cup_{\alpha \in \mathcal{A}} T_\alpha$ such that the following two conditions hold:

- (1) $\Gamma(t_1, t_2) \neq +\infty$ if and only if there is $\alpha \in \mathcal{A}$ such that $t_1 \in T_\alpha$ and $t_2 \in T_\alpha$;
- (2) for every $\alpha \in \mathcal{A}$, the restriction of Γ to $T_\alpha \times T_\alpha$ is negative definite in the usual sense.

Note that the above conditions show that Γ determines the decomposition $T = \cup_{\alpha \in \mathcal{A}} T_\alpha$ uniquely. The next simple proposition establishes a link between positive-definite and negative-definite kernels.

Proposition 2.1. *For each $n \in \mathbb{N}$, let $\{X_n(t), t \in T\}$ be a zero-mean, unit-variance Gaussian process with covariance function $r_n(t_1, t_2) := \mathbb{E}[X_n(t_1)X_n(t_2)]$. Suppose that there exists a sequence z_n with $\lim_{n \rightarrow \infty} z_n = +\infty$ such that for all $t_1, t_2 \in T$,*

$$(5) \quad \Gamma(t_1, t_2) := \lim_{n \rightarrow \infty} z_n(1 - r_n(t_1, t_2)) \in [0, \infty].$$

Then the kernel Γ is negative definite in the extended sense.

Proof. First we show that if for some $t_1, t_2, t_3 \in T$, both $\Gamma(t_1, t_2)$ and $\Gamma(t_2, t_3)$ are finite, then $\Gamma(t_1, t_3)$ is finite too. If the process $\{X_n(t), t \in T\}$ is defined on some probability space $(\Omega, \mathbb{P})$, then we can define an embedding of T into $L^2(\Omega)$ by $t \mapsto X_n(t)$. This embedding induces a pseudometric d_n on T defined by

$$d_n^2(t_1, t_2) = 2(1 - r_n(t_1, t_2)).$$

Now suppose that $\Gamma(t_1, t_2)$ and $\Gamma(t_2, t_3)$ are finite, which means that

$$\lim_{n \rightarrow \infty} z_n^{1/2} d_n(t_1, t_2) < \infty, \quad \lim_{n \rightarrow \infty} z_n^{1/2} d_n(t_2, t_3) < \infty.$$

By the triangle inequality, $d_n(t_1, t_3) \leq d_n(t_1, t_2) + d_n(t_2, t_3)$. It follows that

$$\limsup_{n \rightarrow \infty} z_n^{1/2} d_n(t_1, t_3) < \infty.$$

This implies that $\Gamma(t_1, t_3)$ is finite too.

Now we define an equivalence relation “ $\sim$ ” on T by declaring $t_1 \sim t_2$ if and only if $\Gamma(t_1, t_2) \neq +\infty$. Let $T = \cup_{\alpha \in \mathcal{A}} T_\alpha$ be the decomposition of T into equivalence classes induced by the relation “ $\sim$ ”. To complete the proof, we need to show that for every $\alpha \in \mathcal{A}$, the restriction of Γ to $T_\alpha \times T_\alpha$ is negative definite in the usual sense. Take some $k \in \mathbb{N}$, $t_1, \dots, t_k \in T_\alpha$ and $a_1, \dots, a_k \in \mathbb{R}$ satisfying $\sum_{i=1}^k a_i = 0$. Then, by (5),

$$\sum_{i=1}^k \sum_{j=1}^k a_i a_j \Gamma(t_i, t_j) = \lim_{n \rightarrow \infty} z_n \sum_{i=1}^k \sum_{j=1}^k a_i a_j (1 - r_n(t_i, t_j)).$$

Since $\sum_{i=1}^k a_i = 0$, we have

$$\sum_{i=1}^k \sum_{j=1}^k a_i a_j \Gamma(t_i, t_j) = - \lim_{n \rightarrow \infty} z_n \sum_{i=1}^k \sum_{j=1}^k a_i a_j r_n(t_i, t_j).$$

Now, the right-hand side is ≤ 0 since the kernel $(t', t'') \mapsto r_n(t', t'')$, being the covariance function of the process X_n , is positive definite. This verifies (4) and completes the proof. $\square$

2.2. A family of Gauss–Poisson point processes. In the next theorem we construct a family of exponential intensity Poisson processes related to each other by Gaussian increments. This construction will be used to define limiting processes for both maxima and minima of independent Gaussian processes.

Theorem 2.1. *Fix $\lambda \in \mathbb{R}$ and a negative-definite kernel Γ on a set T . Let $\{U_i, i \in \mathbb{N}\}$ be a Poisson point process on $\mathbb{R}$ with intensity $e^{-\lambda u} du$, and let W_i , $i \in \mathbb{N}$, be independent copies of any zero-mean Gaussian process $\{W(t), t \in T\}$ with incremental variance Γ . Denote by $\sigma^2(t)$ the variance of $W(t)$. Define a function $\mathcal{U}_i : T \rightarrow \mathbb{R}$ by*

$$(6) \quad \mathcal{U}_i(t) = U_i + W_i(t) - \lambda \sigma^2(t)/2.$$

Then the law of the random family of functions $\{\mathcal{U}_i, i \in \mathbb{N}\}$ depends only on Γ and λ (and does not depend on σ^2).

Remark 2.1. For a given negative-definite kernel Γ , there are many processes W with incremental variance Γ . For example, for every $s \in T$ there is a unique in law Gaussian process $W^{(s)}$ with incremental variance Γ such that additionally, $W^{(s)}(s) = 0$. The covariance function of this process is given by

$$(7) \quad \mathbb{E}[W^{(s)}(t_1)W^{(s)}(t_2)] = \frac{1}{2}(\Gamma(t_1, s) + \Gamma(t_2, s) - \Gamma(t_1, t_2)).$$

Remark 2.2. We consider $\{\mathcal{U}_i, i \in \mathbb{N}\}$ as a Poisson point process on the space $\mathbb{R}^T$ endowed with the product σ -algebra.

Remark 2.3. For every fixed $t \in T$, $\{\mathcal{U}_i(t), i \in \mathbb{N}\}$ is a Poisson point process on $\mathbb{R}$ with intensity $e^{-\lambda u} du$. We may view the map $t \mapsto \{\mathcal{U}_i(t), i \in \mathbb{N}\}$ as a family of dependent Poisson point processes on $\mathbb{R}$ related to each other via Gaussian jumps.

Remark 2.4. If $T = \mathbb{R}^d$ and W is a process with *stationary increments*, the statement of Theorem 2.1 can be deduced from [9]. In this case, the law of the family of functions $\{\mathcal{U}_i, i \in \mathbb{N}\}$ is translation invariant. The proof in the general case will be given in Section 6.

3. MAXIMA OF INDEPENDENT GAUSSIAN PROCESSES

The main result of this section is Theorem 3.2 which solves Problem 1. Before we can state and prove it, we need to define the class of limiting processes in Section 3.1 and to recall a result of Hüsler and Reiss [5] in a slightly generalized form and with a different representation of the limiting process in Section 3.2.

3.1. Construction of limiting processes. Let Γ be a negative-definite kernel on a set T . Let $\{U_i, i \in \mathbb{N}\}$ be a Poisson point process on $\mathbb{R}$ with intensity $e^{-u} du$, and let $W_i, i \in \mathbb{N}$, be independent copies of a zero-mean Gaussian process $\{W(t), t \in T\}$ with incremental variance Γ and variance σ^2 . Then we set

$$(8) \quad M_\Gamma(t) = \max_{i \in \mathbb{N}} (U_i + W_i(t) - \sigma^2(t)/2).$$

Remark 3.1. By Theorem 2.1 with $\lambda = 1$, the law of M_Γ depends only on Γ .

Remark 3.2. A standard calculation with Poisson point processes, see [14], shows that the finite-dimensional distributions of M_Γ are as follows: for every $t_1, \dots, t_k \in T$ and $y_1, \dots, y_k \in \mathbb{R}$, we have

$$(9) \quad \begin{aligned} \mathbb{P}[M_\Gamma(t_1) \leq y_1, \dots, M_\Gamma(t_k) \leq y_k] \\ = \exp \left(- \int_{\mathbb{R}} \mathbb{P}[\exists j : u + W(t_j) - \sigma^2(t_j)/2 > y_j] e^{-u} du \right). \end{aligned}$$

Note that for every fixed $t \in T$, $M_\Gamma(t)$ is distributed according to the unit Gumbel distribution function $e^{-e^{-y}}$.

Remark 3.3. By construction, the process M_Γ is max-stable, see [4]. This means that for every $k \in \mathbb{N}$, the process $\{\max_{i=1, \dots, k} M_\Gamma^{(i)}(t), t \in T\}$, where $M_\Gamma^{(1)}, \dots, M_\Gamma^{(k)}$ are independent copies of M_Γ , has the same law as $\{M_\Gamma(t) + \log k, t \in T\}$.

We need to slightly extend the definition of M_Γ given above. Let Γ be a negative-definite kernel in the *extended sense* on a set T with the corresponding decomposition $T = \cup_{\alpha \in \mathcal{A}} T_\alpha$. Then we denote by $\{M_\Gamma(t), t \in T\}$ the stochastic process with the following properties:

- (1) for every $\alpha \in \mathcal{A}$, the process $\{M_\Gamma(t), t \in T_\alpha\}$ is defined as in (8).
- (2) for $\alpha, \beta \in \mathcal{A}$ with $\alpha \neq \beta$, the processes $\{M_\Gamma(t), t \in T_\alpha\}$ and $\{M_\Gamma(t), t \in T_\beta\}$ are independent.

3.2. Sufficient conditions for convergence. The next theorem is a slight modification of a result of Hüsler and Reiss [5]. Closely related results were obtained by [3], [9]. Let u_n be a sequence such that

$$(10) \quad \sqrt{2\pi}u_n e^{u_n^2/2} \sim n, \quad n \rightarrow \infty.$$

Equivalently, one can take

$$(11) \quad u_n = \sqrt{2 \log n} - \frac{(1/2) \log \log n + \log(2\sqrt{\pi}) + o(1)}{\sqrt{2 \log n}}, \quad n \rightarrow \infty.$$

Theorem 3.1. *For every $n \in \mathbb{N}$, let $X_{1n}, \dots, X_{nn}$ be independent copies of a zero-mean, unit-variance Gaussian process $\{X_n(t), t \in T\}$ with covariance function $r_n(t_1, t_2) := \mathbb{E}[X_n(t_1)X_n(t_2)]$ such that for every $t_1, t_2 \in T$,*

$$(12) \quad \Gamma(t_1, t_2) := \lim_{n \rightarrow \infty} 4 \log n \cdot (1 - r_n(t_1, t_2)) \in [0, \infty].$$

Let $M_n(t) = \max_{i=1, \dots, n} X_{in}(t)$. Then the process $u_n(M_n - u_n)$ converges as $n \rightarrow \infty$ to the process M_Γ .

Remark 3.4. By Proposition 2.1, the kernel Γ is negative definite in the extended sense. Hence, the process M_Γ is well defined.

Remark 3.5. Hüsler and Reiss [5] proved Theorem 3.1 assuming that T is finite and that $\Gamma(t_1, t_2) < \infty$ for all $t_1, t_2 \in T$, but they gave a different expression for the limiting process.

Proof of Theorem 3.1. Suppose first that $\Gamma(t_1, t_2) < \infty$ for all $t_1, t_2 \in T$. We will use a method similar to that of [5] (see also [13]). Let $s \in T$ be arbitrary. It will be convenient to set $u_n(y) = u_n + u_n^{-1}y$ for $y \in \mathbb{R}$. Conditioned on the event $A_n(y) := \{X_n(s) = u_n(y)\}$, we have

$$(13) \quad \mathbb{E}[u_n(X_n(t) - u_n) | A_n(y)] = u_n(u_n(y)r_n(t, s) - u_n) = yr_n(t, s) + u_n^2(r_n(t, s) - 1).$$

Taking into account (11) and (12), we obtain

$$(14) \quad \lim_{n \rightarrow \infty} \mathbb{E}[u_n(X_n(t) - u_n) | A_n(y)] = y - \frac{1}{2}\Gamma(t, s).$$

Further, by the well-known formula for the conditional covariance,

$$(15) \quad \text{Cov}[u_n(X_n(t_1) - u_n), u_n(X_n(t_2) - u_n) | A_n(y)] = u_n^2(r_n(t_1, t_2) - r_n(t_1, s)r_n(t_2, s)).$$

Together with (11) and (12) this implies that

$$(16) \quad \lim_{n \rightarrow \infty} \text{Cov}[u_n(X_n(t_1) - u_n), u_n(X_n(t_2) - u_n) | A_n(y)] = \frac{1}{2}(\Gamma(t_1, s) + \Gamma(t_2, s) - \Gamma(t_1, t_2)).$$

It follows from (14) and (16) that for every fixed $y \in \mathbb{R}$, we have the following convergence of conditional processes

$$(17) \quad \{u_n(X_n(t) - u_n) | A_n(y), t \in T\} \xrightarrow{\mathcal{D}} \{y + W^{(s)}(t) - \frac{1}{2}\Gamma(t, s), t \in T\}, \quad n \rightarrow \infty.$$

We are going to compute the finite-dimensional distributions of the process $u_n(M_n - u_n)$ in the limit $n \rightarrow \infty$. Fix some $k \in \mathbb{N}$, $t_1, \dots, t_k \in T$, and $y_1, \dots, y_k \in \mathbb{R}$. The density f_n of the random variable $u_n(X_n(s) - u_n)$ is given by

$$f_n(x) = \frac{1}{\sqrt{2\pi}u_n e^{u_n^2/2}} e^{-x} e^{-\frac{x^2}{2u_n^2}}.$$

Conditioning on the event $A_n(y)$, we obtain

$$\begin{aligned} & \mathbb{P}[\exists j : X_n(t_j) > u_n(y_j)] \\ &= \frac{1}{\sqrt{2\pi}u_n e^{u_n^2/2}} \int_{\mathbb{R}} \mathbb{P}[\exists j : u_n(X_n(t_j) - u_n) > y_j | A_n(y)] e^{-y} e^{-\frac{y^2}{2u_n^2}} dy. \end{aligned}$$

It follows from (10), (17), and a standard argument justifying the use of the dominated convergence theorem that

$$(18) \quad \lim_{n \rightarrow \infty} n \mathbb{P}[\exists j : X_n(t_j) > u_n(y_j)] = \int_{\mathbb{R}} \mathbb{P} \left[\exists j : y + W^{(s)}(t_j) - \frac{1}{2}\Gamma(t_j, s) > y_j \right] e^{-y} dy.$$

By the definition of M_n , we have

$$(19) \quad \mathbb{P}[\forall j : u_n(M_n(t_j) - u_n) \leq y_j] = (1 - \mathbb{P}[\exists j : X_n(t_j) > u_n(y_j)])^n.$$

It follows from (18) and (19) that

$$\begin{aligned} (20) \quad & \lim_{n \rightarrow \infty} \mathbb{P}[\forall j : u_n(M_n(t_j) - u_n) \leq y_j] \\ &= \exp \left(- \int_{\mathbb{R}} \mathbb{P} \left[\exists j : y + W^{(s)}(t_j) - \frac{1}{2}\Gamma(t_j, s) > y_j \right] e^{-y} dy \right). \end{aligned}$$

Comparing this with (9), we obtain the statement of the theorem under the restriction that $\Gamma(t_1, t_2) < \infty$ for all $t_1, t_2 \in T$. (Note that by Remark 3.1, we can take $W = W^{(s)}$ on the right-hand side of (9).)

To prove the general case, let Γ be a negative-definite kernel in the extended sense, with the corresponding decomposition $T = \cup_{\alpha \in A} T_\alpha$. Take $t_1, \dots, t_k \in T$ and $y_1, \dots, y_k \in \mathbb{R}$. Let I_α be the set of all $i = 1, \dots, k$ with the property that $t_i \in T_\alpha$. By the Bonferroni inequality,

$$(21) \quad S_n^{(1)} - S_n^{(2)} \leq \mathbb{P}[\exists j : X_n(t_j) > u_n(y_j)] \leq S_n^{(1)},$$

where

$$(22) \quad S_n^{(1)} = \sum_{\alpha \in A} \mathbb{P}[\exists j \in I_\alpha : X_n(t_j) > u_n(y_j)],$$

$$(23) \quad S_n^{(2)} = \sum_{\substack{i', i''=1, \dots, k \\ i' \sim i''}} \mathbb{P}[X_n(t_{i'}) > u_n(y_{i'}), X_n(t_{i''}) > u_n(y_{i''})].$$

(We write $i' \sim i''$ if i' and i'' are contained in the same I_α). For every $\alpha \in A$, choose $s_\alpha \in T_\alpha$ arbitrarily. On the one hand, the above proof, see (18), shows that

for every $\alpha \in A$,

$$\begin{aligned} & \lim_{n \rightarrow \infty} n \mathbb{P}[\exists j \in I_\alpha : X_n(t_j) > u_n(y_j)] \\ &= \int_{\mathbb{R}} \mathbb{P} \left[\exists j \in I_\alpha : y + W^{(s_\alpha)}(t_j) - \frac{1}{2} \Gamma(t_j, s_\alpha) > y_j \right] e^{-y} dy. \end{aligned}$$

On the other hand, it follows from Lemma 3.1 below that we have $\lim_{n \rightarrow \infty} n S_n^{(2)} = 0$. It follows from (21) that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \mathbb{P}[\forall j : M_n(t_j) \leq u_n(y_j)] \\ &= \exp \left(- \sum_{\alpha \in A} \int_{\mathbb{R}} \mathbb{P} \left[\exists j \in I_\alpha : y + W^{(s_\alpha)}(t_j) - \frac{1}{2} \Gamma(t_j, s_\alpha) > y_j \right] e^{-y} dy \right). \end{aligned}$$

This completes the proof since it follows from (9) and Section 3.1 that the right-hand side is equal to $\mathbb{P}[\forall j : M_\Gamma(t_j) \leq y_j]$. $\square$

Lemma 3.1. *Let X_n , $n \in \mathbb{N}$, be a sequence of Gaussian processes as in Theorem 3.1. Suppose that for some $t_1, t_2 \in T$, $\Gamma(t_1, t_2) = \infty$. Then for every $y_1, y_2 \in \mathbb{R}$,*

$$p_n := n \mathbb{P}[X_n(t_i) > u_n + u_n^{-1} y_i, i = 1, 2] \rightarrow 0, \quad n \rightarrow \infty.$$

Proof. Fix $\lambda > 0$. Let $\{X_n^{(\lambda)}(t), t \in \{t_1, t_2\}\}$ be a zero-mean, unit-variance Gaussian vector with covariance $\rho_n := 1 - \lambda / \log n$. Recalling (12) and taking into account the fact that $\Gamma(t_1, t_2) = \infty$, we obtain that $r_n(t_1, t_2) < \rho_n$ for sufficiently large n . By Slepian's comparison lemma, see [10, Corollary 4.2.3],

$$(24) \quad p_n \leq n \mathbb{P}[X_n^{(\lambda)}(t_1) > u_n + u_n^{-1} y_1, X_n^{(\lambda)}(t_2) > u_n + u_n^{-1} y_2].$$

The limit of the right-hand side of (24) as $n \rightarrow \infty$ was computed by [5]. It follows that

$$\limsup_{n \rightarrow \infty} p_n \leq e^{-y_1} + e^{-y_2} - \Phi \left(\lambda + \frac{y_2 - y_1}{2\lambda} \right) e^{-y_1} - \Phi \left(\lambda + \frac{y_1 - y_2}{2\lambda} \right) e^{-y_2},$$

where Φ is the standard normal distribution function. Since the above is true for every $\lambda > 0$, we obtain the statement of the lemma by letting $\lambda \rightarrow \infty$. $\square$

3.3. Necessary and sufficient conditions for convergence. The next theorem solves Problem 1. To exclude trivial cases, we adopt the following definition: A process $\{M(t), t \in T\}$ is called *nondegenerate* if there is $t \in T$ such that $M(t)$ is not a.s. constant.

Theorem 3.2. *For every $n \in \mathbb{N}$, let $X_{1n}, \dots, X_{nn}$ be independent copies of a zero-mean Gaussian process $\{X_n(t), t \in T\}$ with*

$$(25) \quad r_n(t_1, t_2) := \mathbb{E}[X_n(t_1)X_n(t_2)], \quad \sigma_n^2(t) := r_n(t, t) > 0.$$

Define a process $\{M_n(t), t \in T\}$ by $M_n(t) = \max_{i=1, \dots, n} X_{in}(t)$. Then:

- (1) *there exist sequences $a_n > 0$ and b_n such that the process $a_n(M_n - b_n)$ converges as $n \rightarrow \infty$ to a nondegenerate limit M , if and only if, the following two conditions hold:*
 - (a) *the following limit exists in $[0, \infty]$ for every $t_1, t_2 \in T$:*

$$(26) \quad \Gamma(t_1, t_2) := 4 \lim_{n \rightarrow \infty} \log n \cdot \left(1 - \frac{r_n(t_1, t_2)}{\sigma_n(t_1)\sigma_n(t_2)} \right);$$

(b) the following limit exists in $\mathbb{R}$ for every $t_1, t_2 \in T$:

$$(27) \quad \kappa(t_1, t_2) := 2 \lim_{n \rightarrow \infty} \log n \left(1 - \frac{\sigma_n(t_1)}{\sigma_n(t_2)} \right);$$

(2) if Condition 1a holds, then the kernel Γ is negative definite in the extended sense;

(3) if Conditions 1a and 1b are satisfied, then we can choose

$$(28) \quad a_n = u_n / \sigma_n(t_0), \quad b_n = \sigma_n(t_0) u_n,$$

where $t_0 \in T$ is fixed and u_n is as in (11). In this case, the limiting process $\{M(t), t \in T\}$ has the same law as $\{M_\Gamma(t) + \kappa(t_0, t), t \in T\}$, where M_Γ is as in Section 3.1.

Remark 3.6. If we additionally assume that $\sigma_n \equiv 1$, then Condition 1b is trivially fulfilled with $\kappa(t_1, t_2) = 0$.

We will need the following lemma from [9], see Lemma 21 there.

Lemma 3.2. *Let the assumptions of Theorem 3.2 be satisfied and, additionally, $\sigma_n \equiv 1$. Take some $t_1, t_2 \in T$. Then the law of the bivariate random vector $(u_n(M_n(t_1) - u_n), u_n(M_n(t_2) - u_n))$ converges weakly as $n \rightarrow \infty$ if and only if the following limit exists in $[0, \infty]$:*

$$(29) \quad c := \lim_{n \rightarrow \infty} \log n \cdot (1 - r_n(t_1, t_2)).$$

Proof of Theorem 3.2. Suppose that the process $a_n(M_n - b_n)$ converges as $n \rightarrow \infty$ to some nondegenerate limit M . First we prove the existence of the limit in (27). The random variable $\sigma_n^{-1}(t)M_n(t)$ has the distribution of the maximum of n independent standard Gaussian random variables. Thus, the random variable $u_n \sigma_n^{-1}(t)(M_n(t) - \sigma_n(t)u_n)$ converges as $n \rightarrow \infty$ weakly to the Gumbel distribution, see [10, Theorem 1.5.3]. On the other hand, since we assume the process M to be nondegenerate, for some $t \in T$, the random variable $a_n(M_n(t) - b_n)$ has a nondegenerate limiting distribution as $n \rightarrow \infty$. By the convergence to types lemma, see [10, Theorem 1.2.3], there exist constants $a(t) > 0$ and $b(t) \in \mathbb{R}$ such that

$$(30) \quad \lim_{n \rightarrow \infty} \frac{u_n \sigma_n^{-1}(t)}{a_n} = a(t), \quad \lim_{n \rightarrow \infty} a_n(\sigma_n(t)u_n - b_n) = b(t).$$

Inserting $t = t_1$ and $t = t_2$ into the second equation in (30) and taking the difference, we obtain

$$\lim_{n \rightarrow \infty} a_n u_n (\sigma_n(t_1) - \sigma_n(t_2)) = b(t_1) - b(t_2).$$

Thus,

$$(31) \quad \lim_{n \rightarrow \infty} u_n a_n \sigma_n(t_1) \left(1 - \frac{\sigma_n(t_2)}{\sigma_n(t_1)} \right) = b(t_1) - b(t_2).$$

Recall that by (30), $a_n \sigma_n(t_1) \sim u_n / a(t)$, and by (11), $u_n^2 \sim 2 \log n$ as $n \rightarrow \infty$. Applying this to (31), yields Condition 1b.

We prove that Condition 1a is satisfied. Take some $t_1, t_2 \in T$ and assume that $t_1 \neq t_2$, since otherwise, the limit in (26) is evidently 0. Define a unit-variance Gaussian process $\{X'_n(t), t \in T\}$ by $X'_n(t) = X_n(t) / \sigma_n(t)$. Let $M'_n(t) = M_n(t) / \sigma_n(t)$. We claim that the process $u_n(M'_n - u_n)$ converges as $n \rightarrow \infty$ to a

nondegenerate limit. Recall that we assume that the process $a_n(M_n - b_n)$ converges to M . We may write

$$(32) \quad \{u_n(M'_n(t) - u_n), t \in T\} \stackrel{\mathcal{D}}{=} \{u_n \sigma_n^{-1}(t)(M_n(t) - \sigma_n(t)u_n), t \in T\}.$$

It follows from (30) that the process on the right-hand side of (32) converges to $\{a(t)(M(t) - b(t)), t \in T\}$. The covariance function of X'_n is given by

$$(33) \quad r'_n(t_1, t_2) = r_n(t_1, t_2) / \sigma_n(t_1) \sigma_n(t_2).$$

Condition 1a follows then from Lemma 3.2.

The above proves the “only if” statement of part 1. Since the “if” statement of part 1 will follow from part 3, we proceed to the proof of part 2 of the theorem. Recall that $X'_n(t) = X_n(t) / \sigma_n(t)$. In view of (33), we may rewrite (26) as

$$\Gamma(t_1, t_2) = \lim_{n \rightarrow \infty} 4 \log n \cdot (1 - r'_n(t_1, t_2)).$$

Applying Proposition 2.1 to the process X'_n , we obtain the statement of part 2.

We prove part 3 of the theorem. Suppose that Conditions 1a and 1b hold. By Theorem 3.1, the process $u_n(M'_n - u_n)$ converges to M_Γ as $n \rightarrow \infty$. With a_n and b_n as in (28) we have

$$\{a_n(M_n(t) - b_n), t \in T\} \stackrel{\mathcal{D}}{=} \left\{ \frac{\sigma_n(t)}{\sigma_n(t_0)} u_n(M'_n(t) - u_n) - u_n^2 \left(1 - \frac{\sigma_n(t)}{\sigma_n(t_0)} \right), t \in T \right\}.$$

Note that by (27),

$$\lim_{n \rightarrow \infty} \frac{\sigma_n(t)}{\sigma_n(t_0)} = 1, \quad \lim_{n \rightarrow \infty} u_n^2 \left(1 - \frac{\sigma_n(t)}{\sigma_n(t_0)} \right) = \kappa(t_0, t).$$

This completes the proof of part 3 of the theorem. $\square$

4. MINIMA OF INDEPENDENT GAUSSIAN PROCESSES

In this section we solve Problem 2 using a method similar to that used in Section 3.

4.1. Construction of limiting processes. We start by defining the class of limiting processes. Let Γ be a negative-definite kernel on a set T . Let $\{U_i, i \in \mathbb{N}\}$ be a Poisson point process on $\mathbb{R}$ with Lebesgue measure as intensity, and let $W_i, i \in \mathbb{N}$, be independent copies of a zero-mean Gaussian process $\{W(t), t \in T\}$ with incremental variance Γ and variance σ^2 . Then we define a process $\{L_\Gamma(t), t \in T\}$ by

$$(34) \quad L_\Gamma(t) = \min_{i \in \mathbb{N}} |U_i + W_i(t)|.$$

Remark 4.1. By Theorem 2.1 with $\lambda = 0$, the law of L_Γ depends only on Γ .

Remark 4.2. It follows from the basic properties of Poisson point processes that the finite-dimensional distributions of L_Γ are as follows: for every $t_1, \dots, t_k \in T$ and $y_1, \dots, y_k > 0$, we have

$$(35) \quad \mathbb{P}[L_\Gamma(t_1) > y_1, \dots, L_\Gamma(t_k) > y_k] = \exp \left(- \int_{\mathbb{R}} \mathbb{P}[\exists j : |u + W(t_j)| \leq y_j] du \right).$$

For every fixed $t \in T$, $L_\Gamma(t)$ has an exponential distribution with mean $1/2$.

Remark 4.3. The process L_Γ is max-infinitely divisible, which means (see [14] for a definition) that for every $k \in \mathbb{N}$, we can represent L_Γ as a pointwise minimum of k independent identically distributed processes. This follows from the fact that the Poisson point process on $\mathbb{R}$ with unit intensity can be represented as a union of k independent Poisson point processes with constant intensity $1/k$. Note also that the property of max-infinite divisibility is weaker than that of max-stability, cf. Remark 3.3.

Exactly as in Section 3.1, the above definition of the process L_Γ can be extended to the case when Γ is a negative-definite kernel in the *extended sense*.

4.2. Sufficient condition for convergence. The next theorem is an analogue of Theorem 3.1 for minima (in the sense of absolute value) of independent Gaussian processes. Let

$$(36) \quad w_n = (2\pi)^{-1/2}n.$$

Theorem 4.1. *For every $n \in \mathbb{N}$, let $X_{1n}, \dots, X_{nn}$ be independent copies of a zero-mean, unit-variance Gaussian process $\{X_n(t), t \in T\}$ with covariance function $r_n(t_1, t_2) := \mathbb{E}[X_n(t_1)X_n(t_2)]$ such that for every $t_1, t_2 \in T$,*

$$(37) \quad \Gamma(t_1, t_2) := \frac{1}{\pi} \lim_{n \rightarrow \infty} n^2(1 - r_n(t_1, t_2)) \in [0, \infty].$$

Let $L_n(t) = \min_{i=1, \dots, n} |X_{in}(t)|$. Then the process $w_n L_n$ converges as $n \rightarrow \infty$ to the process L_Γ .

Remark 4.4. By Proposition 2.1, the kernel Γ is negative-definite in the extended sense. Hence, the process L_Γ is well-defined.

Proof. The idea of the proof is analogous to that used in the proof of Theorem 3.1. First suppose that $\Gamma(t_1, t_2) < \infty$ for every $t_1, t_2 \in T$. Fix some $s \in T$. Conditioned on the event $A_n(y) := \{X_n(s) = w_n^{-1}y\}$, we have

$$(38) \quad \lim_{n \rightarrow \infty} \mathbb{E}[w_n X_n(t) | A_n(y)] = \lim_{n \rightarrow \infty} y r_n(t, s) = y.$$

Further, for every $t_1, t_2 \in T$,

$$\text{Cov}[w_n X_n(t_1), w_n X_n(t_2) | A_n(y)] = w_n^2(r_n(t_1, t_2) - r_n(t_1, s)r_n(t_2, s)).$$

It follows that

$$(39) \quad \lim_{n \rightarrow \infty} \text{Cov}[w_n X_n(t_1), w_n X_n(t_2) | A_n(y)] = \frac{1}{2}(\Gamma(t_1, s) + \Gamma(t_2, s) - \Gamma(t_1, t_2)).$$

It follows from (38) and (39) that for every $y \in \mathbb{R}$, we have the following convergence of conditioned stochastic processes:

$$(40) \quad \{w_n X_n(t) | A_n(y), t \in T\} \xrightarrow{\mathcal{D}} \{y + W^{(s)}(t), t \in T\}, \quad n \rightarrow \infty,$$

where $W^{(s)}$ is as in Remark 2.1.

Take $k \in \mathbb{N}$, $t_1, \dots, t_k \in T$, and $y_1, \dots, y_k \geq 0$. The density f_n of the random variable $w_n X_n(s)$ is given by

$$f_n(y) = \frac{1}{\sqrt{2\pi}w_n} e^{-\frac{y^2}{2w_n^2}}.$$

Conditioning on the event $A_n(y)$, we get

$$(41) \quad \mathbb{P}[\exists j : w_n |X_n(t_j)| \leq y_j] = \frac{1}{\sqrt{2\pi}w_n} \int_{\mathbb{R}} \mathbb{P}[\exists j : w_n |X_n(t_j)| \leq y_j | A_n(y)] e^{-\frac{y^2}{2w_n^2}} dy.$$

It follows from (40) and (41) that

$$\lim_{n \rightarrow \infty} n \mathbb{P}[\exists j : w_n |X_n(t_j)| \leq y_j] = \int_{\mathbb{R}} \mathbb{P}[\exists j : |y + W^{(s)}(t_j)| \leq y_j] dy.$$

Again, we have omitted the standard justification of the use of the dominated convergence theorem. Recalling that L_n is the minimum of n independent copies of $|X_n|$, we obtain

$$\lim_{n \rightarrow \infty} \mathbb{P}[\forall j : w_n L_n(t_j) > y_j] = \exp \left(- \int_{\mathbb{R}} \mathbb{P}[\exists j : |y + W^{(s)}(t_j)| \leq y_j] dy \right).$$

Comparing this with (35) (we may take $W = W^{(s)}$ there) completes the proof of the theorem under the restriction that $\Gamma(t_1, t_2)$ is finite for all $t_1, t_2 \in T$. The general case can be handled exactly as in the proof of Theorem 3.1 using Lemma 4.1 below. $\square$

Lemma 4.1. *Let X_n , $n \in \mathbb{N}$, be a sequence of Gaussian processes as in Theorem 4.1. Suppose that for some $t_1, t_2 \in T$, $\Gamma(t_1, t_2) = \infty$. Then for every $y_1, y_2 > 0$,*

$$p_n := n \mathbb{P}[w_n |X_n(t_i)| < y_i, i = 1, 2] \rightarrow 0, \quad n \rightarrow \infty.$$

Proof. The density g_n of the bivariate vector $(w_n X_n(t_1), w_n X_n(t_2))$ is given by

$$g_n(z_1, z_2) = \frac{1}{2\pi w_n^2 \sqrt{1 - r_n(t_1, t_2)^2}} \exp \left(- \frac{z_1^2 + z_2^2 - 2r_n(t_1, t_2)z_1 z_2}{2w_n^2(1 - r_n(t_1, t_2)^2)} \right).$$

It follows from (46) together with $\Gamma(t_1, t_2) = \infty$ that the term $\exp(\dots)$ on the right-hand side converges to 1 as $n \rightarrow \infty$. Furthermore, $\lim_{n \rightarrow \infty} n g_n(z_1, z_2) = 0$. Thus,

$$p_n = \int_{-y_1}^{y_1} \int_{-y_2}^{y_2} n g_n(z_1, z_2) dz_1 dz_2 \rightarrow 0, \quad n \rightarrow \infty.$$

This completes the proof of the lemma. $\square$

4.3. Necessary and sufficient conditions. The next theorem solves Problem 2.

Theorem 4.2. *For every $n \in \mathbb{N}$, let $X_{1n}, \dots, X_{nn}$ be independent copies of a zero-mean Gaussian process $\{X_n(t), t \in T\}$ with*

$$(42) \quad r_n(t_1, t_2) := \mathbb{E}[X_n(t_1)X_n(t_2)], \quad \sigma_n^2(t) := r_n(t, t) > 0.$$

Define a process $\{L_n(t), t \in T\}$ by $L_n(t) = \min_{i=1, \dots, n} |X_{in}(t)|$. Then:

- (1) *there exists a sequence a_n such that the process $a_n L_n$ converges as $n \rightarrow \infty$ to some nondegenerate limit L , if and only if, the following conditions hold:*
 - (a) *the following limit exists in $[0, \infty]$ for every $t_1, t_2 \in T$:*

$$(43) \quad \Gamma(t_1, t_2) := \frac{1}{\pi} \lim_{n \rightarrow \infty} n^2 \left(1 - \frac{r_n(t_1, t_2)}{\sigma_n(t_1)\sigma_n(t_2)} \right);$$

- (b) *the following limit exists in $(0, \infty)$ for every $t_1, t_2 \in T$:*

$$(44) \quad \kappa(t_1, t_2) := \lim_{n \rightarrow \infty} \frac{\sigma_n(t_1)}{\sigma_n(t_2)};$$

- (2) *if Condition 1a holds, then the kernel Γ is negative definite in the extended sense;*

(3) if Conditions 1a and 1b are satisfied, then we can choose

$$(45) \quad a_n = (2\pi)^{-1/2} \sigma_n^{-1}(t_0) n$$

for some fixed t_0 . In this case, the limiting process $\{L(t), t \in T\}$ has the same law as $\{\kappa(t, t_0) L_\Gamma(t), t \in T\}$, where L_Γ is as in Section 4.1.

Remark 4.5. Considering a more general normalization $a_n(L_n - b_n)$ will not lead to any essentially new processes by the convergence to types lemma.

We will need the following two lemmas.

Lemma 4.2. *Let the assumptions of Theorem 4.1 be satisfied and, additionally, $\sigma_n \equiv 1$. Take some $t_1, t_2 \in T$. Then the law of the bivariate random vector $(w_n L_n(t_1), w_n L_n(t_2))$ converges weakly as $n \rightarrow \infty$ if and only if the following limit exists in $[0, \infty]$:*

$$(46) \quad c := \lim_{n \rightarrow \infty} n^2(1 - r_n(t_1, t_2)).$$

Proof. The “if” part of the lemma follows from Theorem 4.1. To prove the “only if” part, suppose that the limit on the right-hand side of (46) does not exist, which means that the sequence $n^2(1 - r_n(t_1, t_2))$ has at least two different accumulation points c_1, c_2 in $[0, \infty]$. It follows from Theorem 4.1 that the sequence of vectors $(w_n L_n(t_1), w_n L_n(t_2))$ has at least two weak accumulation points, and these points are different, as it can be seen from the explicit formula (35). This contradiction completes the proof. $\square$

Lemma 4.3. *If Z_i , $i \in \mathbb{N}$, are independent standard normal variables, then the random variable $w_n \min_{i=1, \dots, n} |Z_i|$, where w_n is as in (36), converges as $n \rightarrow \infty$ to the exponential distribution with mean $1/2$.*

Proof. Let $y > 0$. Since the density of Z_1 is equal to $1/\sqrt{2\pi}$ at 0, we have

$$\lim_{n \rightarrow \infty} n \mathbb{P}[|Z_1| \leq w_n^{-1} y] = \lim_{n \rightarrow \infty} \frac{2w_n^{-1} n y}{\sqrt{2\pi}} = 2y.$$

It follows that

$$\lim_{n \rightarrow \infty} \mathbb{P} \left[w_n \min_{i=1, \dots, n} |Z_i| > y \right] = \lim_{n \rightarrow \infty} (1 - \mathbb{P}[|Z_1| \leq w_n^{-1} y])^n = e^{-2y}.$$

This completes the proof. $\square$

Proof of Theorem 4.2. We prove the “only if” statement of part 1 of the theorem. Suppose that the process $a_n L_n$ converges as $n \rightarrow \infty$ to some nondegenerate limit L . First we show that Condition 1b holds. The random variable $\sigma_n^{-1}(t) L_n(t)$ has the distribution of the minimum (in the sense of absolute value) of n independent standard Gaussian random variables. By Lemma 4.3, the random variable $w_n \sigma_n^{-1}(t) L_n(t)$ converges as $n \rightarrow \infty$ weakly to the exponential distribution with mean $1/2$. On the other hand, by the assumption, the random variable $a_n L_n(t)$ has a nondegenerate limiting distribution as $n \rightarrow \infty$. By the convergence to types lemma, see [10, Theorem 1.2.3], there exists a constant $a(t) > 0$ such that

$$(47) \quad \lim_{n \rightarrow \infty} \frac{w_n \sigma_n^{-1}(t)}{a_n} = a(t).$$

Using this for $t = t_1$ and $t = t_2$ and taking the quotient, we obtain (44).

We prove that Condition 1a holds. Take $t_1, t_2 \in T$ with $t_1 \neq t_2$ (otherwise, the limit in (43) is 0). Define a unit-variance Gaussian process $\{X'_n(t), t \in T\}$ by $X'_n(t) = X_n(t)/\sigma_n(t)$, and let $L'_n(t) = L_n(t)/\sigma_n(t)$. We show that the process $w_n L'_n$ converges as $n \rightarrow \infty$. We have

$$\{w_n L'_n(t), t \in T\} \stackrel{\mathcal{D}}{=} \left\{ \frac{w_n \sigma_n^{-1}(t)}{a_n} a_n L_n(t), t \in T \right\}.$$

It follows from (47) that the process $w_n L'_n$ converges as $n \rightarrow \infty$ to $\{a(t)L(t), t \in T\}$. Condition 1a follows then from Lemma 4.2.

To prove part 2 of the theorem, note that if Condition 1a is satisfied, then r'_n , the covariance function of X'_n , satisfies

$$\lim_{n \rightarrow \infty} n^2 \cdot (1 - r'_n(t_1, t_2)) = \pi \Gamma(t_1, t_2).$$

Applying Proposition 2.1 to the process X'_n , we obtain the statement of part 2.

We prove part 3. If Condition 1a is satisfied, then by Theorem 4.1, the process $w_n L'_n$ converges to L_Γ . If a_n is chosen as in (45), then

$$\{a_n L_n(t), t \in T\} \stackrel{\mathcal{D}}{=} \left\{ \frac{\sigma_n(t)}{\sigma_n(t_0)} w_n L'_n(t), t \in T \right\}.$$

It follows from Condition 1b that the process on the right-hand side converges as $n \rightarrow \infty$ to the process $\{\kappa(t, t_0)L_\Gamma(t), t \in T\}$. This completes the proof. $\square$

5. EXAMPLES AND APPLICATIONS

5.1. Maxima of independent fractional Brownian motions. Recall that a zero-mean Gaussian process $\{B(t), t \in \mathbb{R}\}$ is called fractional Brownian motion with index α if

$$(48) \quad \text{Cov}(B(t_1), B(t_2)) = \frac{1}{2} (|t_1|^\alpha + |t_2|^\alpha - |t_1 - t_2|^\alpha).$$

The next theorem describes the limiting processes for maxima of independent fractional Brownian motions. For the usual Brownian motion (part 2 of the theorem), it reduces to a result of Brown and Resnick [3] (whose proof was based on the Markov property of the Brownian motion). On the other hand, maxima of independent *stationary* Gaussian processes were studied by [9]. We call a process $\{M(t), t \in T\}$ *nontrivial* if it is nondegenerate (i.e., $M(t)$ is not a.s. constant for at least one $t \in T$), and, additionally, there are $t_1, t_2 \in T$ such that $M(t_1)$ and $M(t_2)$ are not a.s. equal.

Theorem 5.1. *Let B_i , $i \in \mathbb{N}$, be independent copies of a fractional Brownian motion $\{B(t), t \geq 0\}$ with index $\alpha \in (0, 2]$. Fix $t_0 > 0$ and, with $s_n > 0$ to be specified later, define*

$$M_n(t) = \max_{i=1, \dots, n} B_i(t_0 + s_n t).$$

- (1) *Suppose that $\alpha \in (0, 1)$. Define u_n as in (11), and let $s_n = t_0/(2 \log n)^{1/\alpha}$. Then the process $u_n(M_n - u_n)$ converges as $n \rightarrow \infty$ to the process $\{M_\Gamma(t), t \in \mathbb{R}\}$, where $\Gamma(t_1, t_2) = |t_1 - t_2|^\alpha$.*
- (2) *Suppose that $\alpha = 1$. Define u_n as in (11), and let $s_n = t_0/(2 \log n)$. Then the process $u_n(M_n - u_n)$ converges as $n \rightarrow \infty$ to the process $\{M_\Gamma(t) + t/2, t \in \mathbb{R}\}$, where $\Gamma(t_1, t_2) = |t_1 - t_2|$.*

- (3) Suppose that $\alpha \in (1, 2]$. Then there exist no sequences a_n, b_n , and $s_n > 0$ for which the process $a_n(M_n - b_n)$ converges to a nontrivial limiting process.

Remark 5.1. An additional rescaling represented by the sequence s_n is needed to obtain a nontrivial limit, cf. [3, 9].

Remark 5.2. Using methods of [9], it can be shown that the convergence in parts 1 and 2 is weak on $C([-A, A])$, the space of continuous functions on $[-A, A]$, for every $A > 0$.

Proof of Theorem 5.1. We are going to apply Theorem 3.2 with $X_n(t) = B(t_0 + s_n t)$. First, let s_n be an arbitrary sequence with $s_n \rightarrow 0$. The variance $\sigma_n^2(t) := \text{Var } X_n(t)$ satisfies, by (48),

$$(49) \quad \sigma_n(t) = (t_0 + s_n t)^{\alpha/2} = t_0^{\alpha/2} \left(1 + \frac{\alpha t}{2t_0} s_n + o(s_n) \right), \quad n \rightarrow \infty.$$

Again using (48), we obtain for $r_n(t_1, t_2) := \text{Cov}(X_n(t_1), X_n(t_2))$, the covariance function of X_n ,

$$(50) \quad r_n(t_1, t_2) = t_0^\alpha \left(1 + \frac{\alpha(t_1 + t_2)}{2t_0} s_n + o(s_n) \right) - \frac{1}{2} s_n^\alpha |t_1 - t_2|^\alpha, \quad n \rightarrow \infty.$$

It follows from (49) that for every $t_1, t_2 \in \mathbb{R}$,

$$(51) \quad 1 - \frac{\sigma_n(t_1)}{\sigma_n(t_2)} = \frac{\alpha(t_2 - t_1)}{2t_0} s_n + o(s_n), \quad n \rightarrow \infty.$$

We prove part 1 of the theorem. Suppose that $\alpha \in (0, 1)$ and recall that $s_n = t_0/(2 \log n)^{1/\alpha}$. Then $s_n = o(s_n^\alpha)$ as $n \rightarrow \infty$, and an easy calculation based on (49) and (50) yields

$$(52) \quad 1 - \frac{r_n(t_1, t_2)}{\sigma_n(t_1)\sigma_n(t_2)} = \frac{|t_1 - t_2|^\alpha}{2t_0^\alpha} s_n^\alpha + o(s_n^\alpha), \quad n \rightarrow \infty.$$

Hence, Condition 1a of Theorem 3.2 is fulfilled with $\Gamma(t_1, t_2) = |t_1 - t_2|^\alpha$. Further, it follows from (51) that Condition 1b of Theorem 3.2 is fulfilled with $\kappa(t_1, t_2) = 0$. The statement of part 1 follows.

We prove part 2 of the theorem. Suppose that $\alpha = 1$. Then $s_n^\alpha = s_n$ and thus, a calculation based on (49) and (50) yields

$$(53) \quad 1 - \frac{r_n(t_1, t_2)}{\sigma_n(t_1)\sigma_n(t_2)} = \frac{|t_1 - t_2|}{2t_0} s_n + o(s_n), \quad n \rightarrow \infty.$$

Hence, Condition 1a of Theorem 3.2 is fulfilled with $\Gamma(t_1, t_2) = |t_1 - t_2|$. Further, it follows from (51) that Condition 1b of Theorem 3.2 is fulfilled with $\kappa(t_1, t_2) = (t_2 - t_1)/2$. The statement of part 2 follows.

Finally, we prove part 3 of the theorem. Assume that $\alpha \in (1, 2]$ and that for some a_n, b_n , and $s_n > 0$, the process $a_n(M_n - b_n)$ converges to some nontrivial limit M . First of all, note that $\lim_{n \rightarrow \infty} s_n = 0$, since otherwise, Condition 1b of Theorem 3.2 is not fulfilled. We have $s_n^\alpha = o(s_n)$ as $n \rightarrow \infty$, and an easy calculation based on (49) and (50) shows that

$$(54) \quad 1 - \frac{r_n(t_1, t_2)}{\sigma_n(t_1)\sigma_n(t_2)} = o(s_n), \quad n \rightarrow \infty.$$

It follows from (51) that in order that Condition 1b of Theorem 3.2 be fulfilled, we need that $\lim_{n \rightarrow \infty} s_n \log n$ exists finitely. This implies that $s_n = O(1/\log n)$.

as $n \rightarrow \infty$. It follows from (54) that in Condition 1a of Theorem 3.2, we have $\gamma(t_1, t_2) = 0$. Thus, $M(t_1) = M(t_2)$ for all $t_1, t_2 \in T$, which means that the limiting process M is trivial. $\square$

5.2. Minima of independent fractional Brownian motions. The next theorem describes possible limits for minima of independent fractional Brownian motions. In contrast with Theorem 5.1, there is no change in behavior at $\alpha = 1$ here.

Theorem 5.2. *Let B_i , $i \in \mathbb{N}$, be independent copies of a fractional Brownian motion $\{B(t), t \geq 0\}$ with index $\alpha \in (0, 2]$. Fix $t_0 > 0$, and, with $s_n = t_0(2\pi/n)^{1/\alpha}$, define*

$$L_n(t) = \min_{i=1, \dots, n} |B_i(t_0 + s_n t)|.$$

Then the process $(2\pi)^{-1/2} n L_n$ converges as $n \rightarrow \infty$ to the process $\{L_\Gamma(t), t \in \mathbb{R}\}$, where $\Gamma(t_1, t_2) = |t_1 - t_2|^\alpha$.

Remark 5.3. It can be shown that the convergence is weak on the space $C([-A, A])$ for every $A > 0$.

Remark 5.4. It is possible to obtain a generalization of Theorem 4.1, Theorem 4.2 and Theorem 5.2 to χ^2 -processes. Recall that a d -dimensional χ^2 -process is defined as the Euclidian norm of an $\mathbb{R}^d$ -valued process with i.i.d. Gaussian components. This, in particular, recovers the results of [11, 12], where minima of independent Bessel processes were studied. Note that the proofs of [11, 12] are based on the Markov property of Bessel processes. Global minima of stationary χ^2 -processes were studied by [1].

Proof. We will apply Theorem 4.2 to the process $X_n(t) = B(t_0 + s_n t)$. Since $\lim_{n \rightarrow \infty} s_n = 0$, we have, for every $t \in \mathbb{R}$,

$$\lim_{n \rightarrow \infty} \sigma_n(t) = |t_0|^{\alpha/2}.$$

This implies that Condition 1b of Theorem 4.2 holds with $\kappa(t_1, t_2) = 1$. As in the proof of Theorem 5.1, the variance and the covariance of X_n satisfy (49) and (50). Hence,

$$1 - \frac{r_n(t_1, t_2)}{\sigma_n(t_1)\sigma_n(t_2)} = \frac{|t_1 - t_2|^\alpha}{2t_0^\alpha} s_n^\alpha + o(s_n^\alpha), \quad n \rightarrow \infty.$$

It follows that Condition 1a of Theorem 4.2 holds with $\Gamma(t_1, t_2) = |t_1 - t_2|^\alpha$. $\square$

5.3. α -stable fields indexed by metric spaces. Let (T, d) be a pseudo-metric space admitting an isometric embedding into a Hilbert space. We are going to use Theorem 2.1 to construct a natural α -stable process on T whose dependence structure is determined by the metric d . Recall that the kernel Γ defined by $\Gamma(t_1, t_2) := d^2(t_1, t_2)$ is negative definite. Let $\{U_i, i \in \mathbb{N}\}$ be a Poisson point process on $\mathbb{R}$ with intensity $e^{-u} du$ and let $W_i, i \in \mathbb{N}$, be independent copies of any zero-mean Gaussian process $\{W(t), t \in T\}$ with incremental variance Γ . Let also $\sigma^2(t) = \text{Var } W(t)$.

Theorem 5.3. *For $\alpha \in (0, 2)$, define a process $\{S_{\Gamma, \alpha}(t), t \in T\}$ by*

$$(55) \quad S_{\Gamma, \alpha}(t) = \sum_{i \in \mathbb{N}} \left(e^{\alpha(U_i + W_i(t) - \sigma^2(t)/2)} - b_i^{(\alpha)} \right),$$

where

$$b_i^{(\alpha)} = \begin{cases} 0, & 0 < \alpha < 1, \\ \int_{1/i}^{1/(i-1)} x^{-2} \sin x dx, & \alpha = 1, \\ \frac{\alpha}{\alpha-1} (i^{\frac{\alpha}{\alpha-1}} - (i-1)^{\frac{\alpha}{\alpha-1}}), & 1 < \alpha < 2. \end{cases}$$

Then the process $S_{\Gamma, \alpha}$ is an α -stable process totally skewed to the right, and the law of $S_{\Gamma, \alpha}$ depends only on Γ and α (and does not depend on σ^2).

Proof. The α -stability of $S_{\Gamma, \alpha}$ follows from the series representation of multivariate α -stable distributions, see [15, Theorem 3.10.1]. The fact that $S_{\Gamma, \alpha}$ depends only on Γ and α follows from Theorem 2.1, since $S_{\Gamma, \alpha}$ can be viewed as a functional of the random family of functions $\{\mathcal{U}_i, i \in \mathbb{N}\}$. $\square$

Remark 5.5. The above approach can be modified to construct *symmetric* α -stable processes. To this end, supply the summands on the right-hand side of (55) with random signs.

Remark 5.6. For two classes of α -stable processes indexed by *positive-definite* kernels, sub-Gaussian and harmonizable processes, see [15]. Examples of α -stable processes indexed by certain metric spaces (which are different from ours) were given by [7]. In a particular case of stationary processes on $\mathbb{R}^d$, the possibility to construct processes as in Theorem 5.3 was mentioned in [8].

Example 5.1. Let $T = \mathbb{S}^d$ be the unit d -dimensional sphere and denote by ρ the geodesic distance on $\mathbb{S}^d$. It is known, see e.g. [6], that the kernel $\Gamma(t_1, t_2) := \rho^\beta(t_1, t_2)$ is negative definite for $\beta \in (0, 1)$. This leads to a family of max-stable (Section 3.1) and a family of α -stable (Theorem 5.3) processes indexed by $\alpha \in (0, 2)$ and $\beta \in (0, 1)$. Note that the laws of these processes are invariant with respect to the rigid rotations of the sphere. A similar construction is possible on a space of constant *negative* curvature.

6. PROOF OF THEOREM 2.1.

STEP 1. Fix $s \in T$ and let $\{W^{(s)}(t), t \in T\}$ be a zero-mean Gaussian process defined as in Remark 2.1, i.e.

$$(56) \quad W^{(s)}(s) = 0, \quad \mathbb{E}[W^{(s)}(t_1)W^{(s)}(t_2)] = \frac{1}{2}(\Gamma(t_1, s) + \Gamma(t_2, s) - \Gamma(t_1, t_2)).$$

Let $\{\mathcal{U}_i^{(s)}, i \in \mathbb{N}\}$ be the random collection of functions constructed as in Theorem 2.1 with $W = W^{(s)}$. More precisely, we define $\mathcal{U}_i^{(s)} : T \rightarrow \mathbb{R}$ by

$$(57) \quad \mathcal{U}_i^{(s)}(t) = U_i + W_i^{(s)}(t) - \lambda\Gamma(t, s)/2.$$

In Step 1 of the proof we are going to show that the law of the random family of functions $\{\mathcal{U}_i^{(s)}, i \in \mathbb{N}\}$ does not depend on the choice of $s \in T$. To this end, we will modify an argument from [9].

Take some $t_1, \dots, t_k \in T$. For a set $B \subset \mathbb{R}^k$ and $x \in \mathbb{R}$, denote by $B + x$ the diagonally shifted set $B + (x, x, \dots, x)$. Let $\mathbb{P}_{t_1, \dots, t_k}^{(s)}$ be the law of the random vector

$$(58) \quad (W^{(s)}(t_1) - \lambda\Gamma(t_1, s)/2, \dots, W^{(s)}(t_k) - \lambda\Gamma(t_k, s)/2),$$

considered as a probability measure on $\mathbb{R}^k$. Then we can view the random collection of points

$$\{(\mathcal{U}_i^{(s)}(t_1), \dots, \mathcal{U}_i^{(s)}(t_k)), i \in \mathbb{N}\}$$

as a Poisson point process on $\mathbb{R}^k$ whose intensity $\Lambda_{t_1, \dots, t_k}^{(s)}$ is given by

$$(59) \quad \Lambda_{t_1, \dots, t_k}^{(s)}(B) = \int_{\mathbb{R}} e^{\lambda x} \mathbb{P}_{t_1, \dots, t_k}^{(s)}(B + x) dx, \quad B \in \mathcal{B}(\mathbb{R}^k).$$

Here, $\mathcal{B}(\mathbb{R}^k)$ is the σ -algebra of Borel subsets of $\mathbb{R}^k$. Consider a measure $\mu_{t_1, \dots, t_k}^{(s)}$ on the $(k-1)$ -dimensional hyperplane $\{(x_i)_{i=1}^k \in \mathbb{R}^k : x_1 = 0\}$ in $\mathbb{R}^k$, defined for $A \in \mathcal{B}(\mathbb{R}^k)$ by

$$(60) \quad \mu_{t_1, \dots, t_k}^{(s)}(A) = \int_{\mathbb{R}^k} e^{\lambda y_1} 1_A(0, y_2 - y_1, \dots, y_k - y_1) d\mathbb{P}_{t_1, \dots, t_k}^{(s)}(y_1, \dots, y_k).$$

Note that $\mu_{t_1, \dots, t_k}^{(s)}(\mathbb{R}^k) = \mathbb{E} e^{\lambda W^{(s)}(t_1) - \lambda^2 \Gamma(t_1, s)/2} = 1$, and therefore, $\mu_{t_1, \dots, t_k}^{(s)}$ is a probability measure. As in [9], we can deduce from (59) that

$$(61) \quad \Lambda_{t_1, \dots, t_k}^{(s)}(B) = \int_{\mathbb{R}} e^{\lambda z} \mu_{t_1, \dots, t_k}^{(s)}(B + z) dz, \quad B \in \mathcal{B}(\mathbb{R}^k).$$

Let $\psi_{t_1, \dots, t_k}^{(s)}(u_1, \dots, u_k) = \int_{\mathbb{R}^k} e^{\sum_{i=1}^k u_i y_i} d\mu_{t_1, \dots, t_k}^{(s)}(y_1, \dots, y_k)$ be the Laplace transform of the measure $\mu_{t_1, \dots, t_k}^{(s)}$. It follows from (60) that

$$(62) \quad \begin{aligned} \psi_{t_1, \dots, t_k}^{(s)}(u_1, \dots, u_k) &= \int_{\mathbb{R}^k} e^{\lambda y_1} e^{u_2(y_2 - y_1) + \dots + u_k(y_k - y_1)} d\mathbb{P}_{t_1, \dots, t_k}^{(s)}(y_1, \dots, y_k) \\ &= \varphi_{t_1, \dots, t_k}^{(s)} \left(\lambda - \sum_{i=2}^k u_i, u_2, \dots, u_k \right), \end{aligned}$$

where $\varphi_{t_1, \dots, t_k}^{(s)} = \int_{\mathbb{R}^k} e^{\sum_{i=1}^k u_i y_i} d\mathbb{P}_{t_1, \dots, t_k}^{(s)}(y_1, \dots, y_k)$ is the Laplace transform of the measure $\mathbb{P}_{t_1, \dots, t_k}^{(s)}$. Recall that the measure $\mathbb{P}_{t_1, \dots, t_k}^{(s)}$ is Gaussian with known covariance and expectation, see (58). Hence,

$$(63) \quad \begin{aligned} \varphi_{t_1, \dots, t_k}^{(s)}(u_1, \dots, u_k) &= \exp \left(-\frac{\lambda}{2} \sum_{i=1}^k \Gamma(t_i, s) u_i + \frac{1}{2} \sum_{i,j=1}^k (\Gamma(t_i, s) + \Gamma(t_j, s) - \Gamma(t_i, t_j)) u_i u_j \right). \end{aligned}$$

An elementary calculation based on (62) and (63) shows that

$$\begin{aligned} \psi_{t_1, \dots, t_k}^{(s)}(u_1, \dots, u_k) &= \exp \left(-\frac{\lambda}{2} \sum_{i=2}^k \Gamma(t_i, t_1) u_i + \frac{1}{2} \sum_{i,j=2}^k (\Gamma(t_i, t_1) + \Gamma(t_j, t_1) - \Gamma(t_i, t_j)) u_i u_j \right). \end{aligned}$$

It follows that the measure $\mu_{t_1, \dots, t_k}^{(s)}$ does not depend on $s \in T$. By (61), this implies that the measure $\Lambda_{t_1, \dots, t_k}^{(s)}$ does not depend on s . Hence, the law of the random family of functions $\{\mathcal{U}_i^{(s)}, i \in \mathbb{N}\}$ does not depend on the choice of $s \in T$.

Remark 6.1. An alternative way to prove the result of Step 1 is as follows. Let $\{X_n(t), t \in T\}$ be a zero-mean Gaussian process whose covariance function is given by

$$r_n(t_1, t_2) = e^{-\Gamma(t_1, t_2)/(4 \log n)}, \quad t_1, t_2 \in T.$$

Note that r_n is indeed a valid covariance function by Schoenberg's theorem, see [2]. The sequence X_n satisfies the assumptions of Theorem 3.1. Taking some $s \in T$ and following the proof of Theorem 3.1, we arrive at (20). Since the left-hand side of (20) does not depend on s , it follows that the right-hand side of (20) does not depend on s either. This implies the statement of Step 1.

STEP 2. Now let $\{W(t), t \in T\}$ be a general zero-mean Gaussian process with incremental variance Γ and variance σ^2 . Our aim is to show that the law of the random family of functions $\{\mathcal{U}_i, i \in \mathbb{N}\}$, where $\mathcal{U}_i$ is defined as in Theorem 2.1, i.e.

$$(64) \quad \mathcal{U}_i(t) = U_i + W_i(t) - \lambda\sigma^2(t)/2,$$

does not depend on σ^2 .

Let $\bar{T}$ be an extension of the set T by a point τ_0 , i.e. $\bar{T} = T \cup \{\tau_0\}$. We extend the kernel Γ to the set $\bar{T}$ as follows: let $\bar{\Gamma}(t_1, t_2) = \Gamma(t_1, t_2)$ if $t_1, t_2 \in T$; $\bar{\Gamma}(\tau_0, t) = \bar{\Gamma}(t, \tau_0) = \sigma^2(t)$; and $\bar{\Gamma}(\tau_0, \tau_0) = 0$. Then $\bar{\Gamma}$ is a negative-definite kernel on $\bar{T}$. To see this, note that $\bar{\Gamma}$ is the incremental variance of the Gaussian process $\{\bar{W}^{(\tau_0)}(t), t \in \bar{T}\}$ defined by $\bar{W}^{(\tau_0)}(\tau_0) = 0$, and $\bar{W}^{(\tau_0)}(t) = W(t)$ for $t \in T$.

Let $s \in \bar{T}$ be arbitrary. Then there is a unique zero-mean Gaussian process $\{\bar{W}^{(s)}(t), t \in \bar{T}\}$ with incremental variance $\bar{\Gamma}$ and $\bar{W}^{(s)}(s) = 0$, see Remark 2.1. Step 1 of the proof implies that the law of the random family of functions $\{\bar{\mathcal{U}}_i^{(s)}, i \in \mathbb{N}\}$, where $\bar{\mathcal{U}}_i^{(s)} : \bar{T} \rightarrow \mathbb{R}$ is defined by

$$(65) \quad \bar{\mathcal{U}}_i^{(s)}(t) = U_i + \bar{W}_i^{(s)}(t) - \lambda\bar{\Gamma}(t, s)/2,$$

does not depend on $s \in T$ (here, $\bar{W}_i^{(s)}, i \in \mathbb{N}$, are independent copies of $\bar{W}^{(s)}$). Note that the restriction of the process $\bar{W}^{(\tau_0)}$ to T has the same law as W , and that for $s \in T$, the restriction of $\bar{W}^{(s)}$ to T has the same law as $W^{(s)}$. Thus, for $s \in T$, the restriction of the family $\{\bar{\mathcal{U}}_i^{(s)}, i \in \mathbb{N}\}$ to T has the same law as the family $\{\mathcal{U}_i^{(s)}, i \in \mathbb{N}\}$ defined in (57), whereas the restriction of the family $\{\bar{\mathcal{U}}_i^{(\tau_0)}, i \in \mathbb{N}\}$ to T has the same law as the family $\{\mathcal{U}_i, i \in \mathbb{N}\}$ defined in (64). It follows that the family $\{\mathcal{U}_i, i \in \mathbb{N}\}$ has the same law as the family $\{\mathcal{U}_i^{(s)}, i \in \mathbb{N}\}$, where $s \in T$ is arbitrary. Since the law of the second family does not depend on σ^2 , we obtain the statement of the theorem. $\square$

ACKNOWLEDGEMENTS

The author is grateful to Martin Schlather for useful remarks.

REFERENCES

- [1] J.M.P. Albin. Minima of H -valued Gaussian processes. *Ann. Probab.*, 24(2): 788–824, 1996.
- [2] C. Berg, J.P.R. Christensen, and P. Ressel. *Harmonic analysis on semigroups. Theory of positive definite and related functions*. Graduate Texts in Mathematics, 100. New York etc.: Springer-Verlag, 1984.
- [3] B.M. Brown and S.I. Resnick. Extreme values of independent stochastic processes. *J. Appl. Probab.*, 14:732–739, 1977.
- [4] L. de Haan. A spectral representation for max-stable processes. *Ann. Probab.*, 12:1194–1204, 1984.

- [5] J. Hüsler and R.-D. Reiss. Maxima of normal random vectors: between independence and complete dependence. *Statist. Probab. Lett.*, 7(4):283–286, 1989.
- [6] J. Istas. Spherical and hyperbolic fractional Brownian motion. *Electron. Commun. Probab.*, 10:254–262, 2005.
- [7] J. Istas. On fractional fields indexed by metric spaces. *Electron. Commun. Probab.*, 11:242–251, 2006.
- [8] Z. Kabluchko. Spectral representations of sum- and max-stable processes. *Extremes*, to appear, 2008.
- [9] Z. Kabluchko, M. Schlather, and L. de Haan. Stationary max-stable fields associated to negative definite functions. *Ann. Probab.*, in press, 2008.
- [10] M.R. Leadbetter, G. Lindgren, and H. Rootzén. *Extremes and related properties of random sequences and processes*. Springer Series in Statistics. New York etc.: Springer-Verlag., 1983.
- [11] M. Penrose. The minimum of a large number of Bessel processes. *J. Lond. Math. Soc., II. Ser.*, 38(3):566–576, 1988.
- [12] M. Penrose. Minima of independent Bessel processes and of distances between Brownian particles. *J. Lond. Math. Soc., II. Ser.*, 43(2):355–366, 1991.
- [13] J. Pickands. Upcrossing probabilities for stationary Gaussian processes. *Trans. Amer. Math. Soc.*, 145:51–73, 1969.
- [14] S.I. Resnick. *Extreme values, regular variation, and point processes*. Applied Probability. Vol. 4. New York etc.: Springer-Verlag., 1987.
- [15] G. Samorodnitsky and M.S. Taqqu. *Stable non-Gaussian random processes: stochastic models with infinite variance*. Stochastic Modeling. New York, NY: Chapman and Hall., 1994.

INSTITUT FÜR MATHEMATISCHE STOCHASTIK, GEORG-AUGUST UNIVERSITÄT GÖTTINGEN, GOLDSCHMIDTSTRASSE 7, D-37077 GÖTTINGEN, GERMANY
E-mail address: `kabluch@math.uni-goettingen.de`